


Palladseite ($\text{Pd}_{17}\text{Se}_{15}$) Structure: A17B15_cP64_207_acfk_eij-001

This structure originally had the label A17B15_cP64_207_acfk_eij. Calls to that address will be redirected here.

Cite this page as: D. Hicks, M. J. Mehl, E. Gossett, C. Toher, O. Levy, R. M. Hanson, G. Hart, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 2*, Comput. Mater. Sci. **161**, S1 (2019). doi: 10.1016/j.commatsci.2018.10.043

<https://aflow.org/p/8LH0>

https://aflow.org/p/A17B15_cP64_207_acfk_eij-001



Prototype	$\text{Pd}_{17}\text{Se}_{15}$
AFLOW prototype label	A17B15_cP64_207_acfk_eij-001
Mineral name	palladseite
ICSD	none
Pearson symbol	cP64
Space group number	207
Space group symbol	$P\bar{4}32$
AFLOW prototype command	<code>aflow --proto=A17B15_cP64_207_acfk_eij-001 --params=a,x₃,x₄,y₅,y₆,x₇,y₇,z₇</code>


Other compounds with this structure

Rh₁₇S₁₅

- (Geller, 1962) determined that $\text{Pd}_{17}\text{Se}_{15}$ could be in space group $Pm\bar{3}m$ #221, $P\bar{4}3m$ #215, or $P432$ #207 (this structure), and finds that $Pm\bar{3}m$ gives the best fit to single-crystal X-ray diffraction pattern, although the parameter fit for the all of the Wyckoff sites did not converge. We therefore present all three structure possibilities.
- We shifted the coordinates of (Geller, 1962) to move the Pd-I atom from the center of the cubic cell, Wyckoff position (1b), to the origin, Wyckoff position (1a).

Simple Cubic primitive vectors

$$\begin{aligned}\mathbf{a}_1 &= a \hat{\mathbf{x}} \\ \mathbf{a}_2 &= a \hat{\mathbf{y}} \\ \mathbf{a}_3 &= a \hat{\mathbf{z}}\end{aligned}$$



Basis vectors

	Lattice coordinates		Cartesian coordinates	Wyckoff position	Atom type
\mathbf{B}_1 =	0	=	0	(1a)	Pd I
\mathbf{B}_2 =	$\frac{1}{2}\mathbf{a}_2 + \frac{1}{2}\mathbf{a}_3$	=	$\frac{1}{2}a\hat{\mathbf{y}} + \frac{1}{2}a\hat{\mathbf{z}}$	(3c)	Pd II
\mathbf{B}_3 =	$\frac{1}{2}\mathbf{a}_1 + \frac{1}{2}\mathbf{a}_3$	=	$\frac{1}{2}a\hat{\mathbf{x}} + \frac{1}{2}a\hat{\mathbf{z}}$	(3c)	Pd II
\mathbf{B}_4 =	$\frac{1}{2}\mathbf{a}_1 + \frac{1}{2}\mathbf{a}_2$	=	$\frac{1}{2}a\hat{\mathbf{x}} + \frac{1}{2}a\hat{\mathbf{y}}$	(3c)	Pd II
\mathbf{B}_5 =	$x_3 \mathbf{a}_1$	=	$ax_3 \hat{\mathbf{x}}$	(6e)	Se I
\mathbf{B}_6 =	$-x_3 \mathbf{a}_1$	=	$-ax_3 \hat{\mathbf{x}}$	(6e)	Se I
\mathbf{B}_7 =	$x_3 \mathbf{a}_2$	=	$ax_3 \hat{\mathbf{y}}$	(6e)	Se I
\mathbf{B}_8 =	$-x_3 \mathbf{a}_2$	=	$-ax_3 \hat{\mathbf{y}}$	(6e)	Se I
\mathbf{B}_9 =	$x_3 \mathbf{a}_3$	=	$ax_3 \hat{\mathbf{z}}$	(6e)	Se I
\mathbf{B}_{10} =	$-x_3 \mathbf{a}_3$	=	$-ax_3 \hat{\mathbf{z}}$	(6e)	Se I
\mathbf{B}_{11} =	$x_4 \mathbf{a}_1 + \frac{1}{2}\mathbf{a}_2 + \frac{1}{2}\mathbf{a}_3$	=	$ax_4 \hat{\mathbf{x}} + \frac{1}{2}a\hat{\mathbf{y}} + \frac{1}{2}a\hat{\mathbf{z}}$	(6f)	Pd III
\mathbf{B}_{12} =	$-x_4 \mathbf{a}_1 + \frac{1}{2}\mathbf{a}_2 + \frac{1}{2}\mathbf{a}_3$	=	$-ax_4 \hat{\mathbf{x}} + \frac{1}{2}a\hat{\mathbf{y}} + \frac{1}{2}a\hat{\mathbf{z}}$	(6f)	Pd III
\mathbf{B}_{13} =	$\frac{1}{2}\mathbf{a}_1 + x_4 \mathbf{a}_2 + \frac{1}{2}\mathbf{a}_3$	=	$\frac{1}{2}a\hat{\mathbf{x}} + ax_4 \hat{\mathbf{y}} + \frac{1}{2}a\hat{\mathbf{z}}$	(6f)	Pd III
\mathbf{B}_{14} =	$\frac{1}{2}\mathbf{a}_1 - x_4 \mathbf{a}_2 + \frac{1}{2}\mathbf{a}_3$	=	$\frac{1}{2}a\hat{\mathbf{x}} - ax_4 \hat{\mathbf{y}} + \frac{1}{2}a\hat{\mathbf{z}}$	(6f)	Pd III
\mathbf{B}_{15} =	$\frac{1}{2}\mathbf{a}_1 + \frac{1}{2}\mathbf{a}_2 + x_4 \mathbf{a}_3$	=	$\frac{1}{2}a\hat{\mathbf{x}} + \frac{1}{2}a\hat{\mathbf{y}} + ax_4 \hat{\mathbf{z}}$	(6f)	Pd III
\mathbf{B}_{16} =	$\frac{1}{2}\mathbf{a}_1 + \frac{1}{2}\mathbf{a}_2 - x_4 \mathbf{a}_3$	=	$\frac{1}{2}a\hat{\mathbf{x}} + \frac{1}{2}a\hat{\mathbf{y}} - ax_4 \hat{\mathbf{z}}$	(6f)	Pd III
\mathbf{B}_{17} =	$y_5 \mathbf{a}_2 + y_5 \mathbf{a}_3$	=	$ay_5 \hat{\mathbf{y}} + ay_5 \hat{\mathbf{z}}$	(12i)	Se II
\mathbf{B}_{18} =	$-y_5 \mathbf{a}_2 + y_5 \mathbf{a}_3$	=	$-ay_5 \hat{\mathbf{y}} + ay_5 \hat{\mathbf{z}}$	(12i)	Se II
\mathbf{B}_{19} =	$y_5 \mathbf{a}_2 - y_5 \mathbf{a}_3$	=	$ay_5 \hat{\mathbf{y}} - ay_5 \hat{\mathbf{z}}$	(12i)	Se II
\mathbf{B}_{20} =	$-y_5 \mathbf{a}_2 - y_5 \mathbf{a}_3$	=	$-ay_5 \hat{\mathbf{y}} - ay_5 \hat{\mathbf{z}}$	(12i)	Se II
\mathbf{B}_{21} =	$y_5 \mathbf{a}_1 + y_5 \mathbf{a}_3$	=	$ay_5 \hat{\mathbf{x}} + ay_5 \hat{\mathbf{z}}$	(12i)	Se II
\mathbf{B}_{22} =	$y_5 \mathbf{a}_1 - y_5 \mathbf{a}_3$	=	$ay_5 \hat{\mathbf{x}} - ay_5 \hat{\mathbf{z}}$	(12i)	Se II
\mathbf{B}_{23} =	$-y_5 \mathbf{a}_1 + y_5 \mathbf{a}_3$	=	$-ay_5 \hat{\mathbf{x}} + ay_5 \hat{\mathbf{z}}$	(12i)	Se II
\mathbf{B}_{24} =	$-y_5 \mathbf{a}_1 - y_5 \mathbf{a}_3$	=	$-ay_5 \hat{\mathbf{x}} - ay_5 \hat{\mathbf{z}}$	(12i)	Se II

References

- [1] S. Geller, *The Crystal Structure of Pd₁₇Se₁₅*, Acta Cryst. **15**, 713–721 (1962), doi:10.1107/S0365110X62001929.

Found in

- [1] D. Barthelmy, *Mineralogy Database* (2012). Palladseite Mineral Data.