


# Ti<sub>3</sub>Al<sub>2</sub>N<sub>2</sub> Structure: A2B4C5\_hP22\_186\_ab\_4b\_a4b-001

Cite this page as: H. Eckert, S. Divilov, A. Zettel, M. J. Mehl, D. Hicks, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 4*. In preparation.

<https://aflow.org/p/7JMK>

[https://aflow.org/p/A2B4C5\\_hP22\\_186\\_ab\\_4b\\_a4b-001](https://aflow.org/p/A2B4C5_hP22_186_ab_4b_a4b-001)




<b>Prototype</b>	Al <sub>2</sub> N <sub>2</sub> Ti <sub>2</sub>
<b>AFLOW prototype label</b>	A2B4C5_hP22_186_ab_4b_a4b-001
<b>ICSD</b>	52643
<b>Pearson symbol</b>	hP22
<b>Space group number</b>	186
<b>Space group symbol</b>	<i>P6<sub>3</sub>mc</i>
<b>AFLOW prototype command</b>	<code>aflow --proto=A2B4C5_hP22_186_ab_4b_a4b-001 --params=a, c/a, z<sub>1</sub>, z<sub>2</sub>, z<sub>3</sub>, z<sub>4</sub>, z<sub>5</sub>, z<sub>6</sub>, z<sub>7</sub>, z<sub>8</sub>, z<sub>9</sub>, z<sub>10</sub>, z<sub>11</sub></code>

- This compound only exists at temperatures near 1573K, where this data was taken.
- (Schuster, 1984) placed this system in the trigonal space group *P31c* #159. The (2a) and (2b) Wyckoff positions for space group #159 are identical to those in the hexagonal space group *P6<sub>3</sub>mc* #186, so we follow (Cenzual, 1991) and place this in the higher symmetry space group.
- The origin of the *z*-axis in space group *P6<sub>3</sub>mc* is arbitrary. We use the origin specified by (Schuster, 1984).
- Most of the Wyckoff positions in this structure are only partially occupied:
  - Sites Al-I and Ti-I are fully occupied.
  - Sites Ni-I, Ni-II, Ti-II, and Ti-III are 80% occupied.
  - Sites Ni-III, Ni-IV, Ti-IV, and Ti-V are 20% occupied.
- This explains the short Ti-N bond lengths shown in the figure – for any given Ti-N pair, only one of the two atoms is actually present.

## Hexagonal primitive vectors

$$\begin{aligned}\mathbf{a}_1 &= \frac{1}{2}a\hat{\mathbf{x}} - \frac{\sqrt{3}}{2}a\hat{\mathbf{y}} \\ \mathbf{a}_2 &= \frac{1}{2}a\hat{\mathbf{x}} + \frac{\sqrt{3}}{2}a\hat{\mathbf{y}} \\ \mathbf{a}_3 &= c\hat{\mathbf{z}}\end{aligned}$$



## Basis vectors

	Lattice coordinates		Cartesian coordinates	Wyckoff position	Atom type
$\mathbf{B}_1$	$= z_1 \mathbf{a}_3$	$=$	$cz_1 \hat{\mathbf{z}}$	(2a)	Al I
$\mathbf{B}_2$	$= (z_1 + \frac{1}{2}) \mathbf{a}_3$	$=$	$c(z_1 + \frac{1}{2}) \hat{\mathbf{z}}$	(2a)	Al I
$\mathbf{B}_3$	$= z_2 \mathbf{a}_3$	$=$	$cz_2 \hat{\mathbf{z}}$	(2a)	Ti I
$\mathbf{B}_4$	$= (z_2 + \frac{1}{2}) \mathbf{a}_3$	$=$	$c(z_2 + \frac{1}{2}) \hat{\mathbf{z}}$	(2a)	Ti I
$\mathbf{B}_5$	$= \frac{1}{3} \mathbf{a}_1 + \frac{2}{3} \mathbf{a}_2 + z_3 \mathbf{a}_3$	$=$	$\frac{1}{2}a\hat{\mathbf{x}} + \frac{\sqrt{3}}{6}a\hat{\mathbf{y}} + cz_3\hat{\mathbf{z}}$	(2b)	Al II
$\mathbf{B}_6$	$= \frac{2}{3} \mathbf{a}_1 + \frac{1}{3} \mathbf{a}_2 + (z_3 + \frac{1}{2}) \mathbf{a}_3$	$=$	$\frac{1}{2}a\hat{\mathbf{x}} - \frac{\sqrt{3}}{6}a\hat{\mathbf{y}} + c(z_3 + \frac{1}{2})\hat{\mathbf{z}}$	(2b)	Al II
$\mathbf{B}_7$	$= \frac{1}{3} \mathbf{a}_1 + \frac{2}{3} \mathbf{a}_2 + z_4 \mathbf{a}_3$	$=$	$\frac{1}{2}a\hat{\mathbf{x}} + \frac{\sqrt{3}}{6}a\hat{\mathbf{y}} + cz_4\hat{\mathbf{z}}$	(2b)	Ni I
$\mathbf{B}_8$	$= \frac{2}{3} \mathbf{a}_1 + \frac{1}{3} \mathbf{a}_2 + (z_4 + \frac{1}{2}) \mathbf{a}_3$	$=$	$\frac{1}{2}a\hat{\mathbf{x}} - \frac{\sqrt{3}}{6}a\hat{\mathbf{y}} + c(z_4 + \frac{1}{2})\hat{\mathbf{z}}$	(2b)	Ni I
$\mathbf{B}_9$	$= \frac{1}{3} \mathbf{a}_1 + \frac{2}{3} \mathbf{a}_2 + z_5 \mathbf{a}_3$	$=$	$\frac{1}{2}a\hat{\mathbf{x}} + \frac{\sqrt{3}}{6}a\hat{\mathbf{y}} + cz_5\hat{\mathbf{z}}$	(2b)	Ni II
$\mathbf{B}_{10}$	$= \frac{2}{3} \mathbf{a}_1 + \frac{1}{3} \mathbf{a}_2 + (z_5 + \frac{1}{2}) \mathbf{a}_3$	$=$	$\frac{1}{2}a\hat{\mathbf{x}} - \frac{\sqrt{3}}{6}a\hat{\mathbf{y}} + c(z_5 + \frac{1}{2})\hat{\mathbf{z}}$	(2b)	Ni II
$\mathbf{B}_{11}$	$= \frac{1}{3} \mathbf{a}_1 + \frac{2}{3} \mathbf{a}_2 + z_6 \mathbf{a}_3$	$=$	$\frac{1}{2}a\hat{\mathbf{x}} + \frac{\sqrt{3}}{6}a\hat{\mathbf{y}} + cz_6\hat{\mathbf{z}}$	(2b)	Ni III
$\mathbf{B}_{12}$	$= \frac{2}{3} \mathbf{a}_1 + \frac{1}{3} \mathbf{a}_2 + (z_6 + \frac{1}{2}) \mathbf{a}_3$	$=$	$\frac{1}{2}a\hat{\mathbf{x}} - \frac{\sqrt{3}}{6}a\hat{\mathbf{y}} + c(z_6 + \frac{1}{2})\hat{\mathbf{z}}$	(2b)	Ni III
$\mathbf{B}_{13}$	$= \frac{1}{3} \mathbf{a}_1 + \frac{2}{3} \mathbf{a}_2 + z_7 \mathbf{a}_3$	$=$	$\frac{1}{2}a\hat{\mathbf{x}} + \frac{\sqrt{3}}{6}a\hat{\mathbf{y}} + cz_7\hat{\mathbf{z}}$	(2b)	Ni IV
$\mathbf{B}_{14}$	$= \frac{2}{3} \mathbf{a}_1 + \frac{1}{3} \mathbf{a}_2 + (z_7 + \frac{1}{2}) \mathbf{a}_3$	$=$	$\frac{1}{2}a\hat{\mathbf{x}} - \frac{\sqrt{3}}{6}a\hat{\mathbf{y}} + c(z_7 + \frac{1}{2})\hat{\mathbf{z}}$	(2b)	Ni IV
$\mathbf{B}_{15}$	$= \frac{1}{3} \mathbf{a}_1 + \frac{2}{3} \mathbf{a}_2 + z_8 \mathbf{a}_3$	$=$	$\frac{1}{2}a\hat{\mathbf{x}} + \frac{\sqrt{3}}{6}a\hat{\mathbf{y}} + cz_8\hat{\mathbf{z}}$	(2b)	Ti II
$\mathbf{B}_{16}$	$= \frac{2}{3} \mathbf{a}_1 + \frac{1}{3} \mathbf{a}_2 + (z_8 + \frac{1}{2}) \mathbf{a}_3$	$=$	$\frac{1}{2}a\hat{\mathbf{x}} - \frac{\sqrt{3}}{6}a\hat{\mathbf{y}} + c(z_8 + \frac{1}{2})\hat{\mathbf{z}}$	(2b)	Ti II
$\mathbf{B}_{17}$	$= \frac{1}{3} \mathbf{a}_1 + \frac{2}{3} \mathbf{a}_2 + z_9 \mathbf{a}_3$	$=$	$\frac{1}{2}a\hat{\mathbf{x}} + \frac{\sqrt{3}}{6}a\hat{\mathbf{y}} + cz_9\hat{\mathbf{z}}$	(2b)	Ti III
$\mathbf{B}_{18}$	$= \frac{2}{3} \mathbf{a}_1 + \frac{1}{3} \mathbf{a}_2 + (z_9 + \frac{1}{2}) \mathbf{a}_3$	$=$	$\frac{1}{2}a\hat{\mathbf{x}} - \frac{\sqrt{3}}{6}a\hat{\mathbf{y}} + c(z_9 + \frac{1}{2})\hat{\mathbf{z}}$	(2b)	Ti III
$\mathbf{B}_{19}$	$= \frac{1}{3} \mathbf{a}_1 + \frac{2}{3} \mathbf{a}_2 + z_{10} \mathbf{a}_3$	$=$	$\frac{1}{2}a\hat{\mathbf{x}} + \frac{\sqrt{3}}{6}a\hat{\mathbf{y}} + cz_{10}\hat{\mathbf{z}}$	(2b)	Ti IV
$\mathbf{B}_{20}$	$= \frac{2}{3} \mathbf{a}_1 + \frac{1}{3} \mathbf{a}_2 + (z_{10} + \frac{1}{2}) \mathbf{a}_3$	$=$	$\frac{1}{2}a\hat{\mathbf{x}} - \frac{\sqrt{3}}{6}a\hat{\mathbf{y}} + c(z_{10} + \frac{1}{2})\hat{\mathbf{z}}$	(2b)	Ti IV
$\mathbf{B}_{21}$	$= \frac{1}{3} \mathbf{a}_1 + \frac{2}{3} \mathbf{a}_2 + z_{11} \mathbf{a}_3$	$=$	$\frac{1}{2}a\hat{\mathbf{x}} + \frac{\sqrt{3}}{6}a\hat{\mathbf{y}} + cz_{11}\hat{\mathbf{z}}$	(2b)	Ti V
$\mathbf{B}_{22}$	$= \frac{2}{3} \mathbf{a}_1 + \frac{1}{3} \mathbf{a}_2 + (z_{11} + \frac{1}{2}) \mathbf{a}_3$	$=$	$\frac{1}{2}a\hat{\mathbf{x}} - \frac{\sqrt{3}}{6}a\hat{\mathbf{y}} + c(z_{11} + \frac{1}{2})\hat{\mathbf{z}}$	(2b)	Ti V

## References

- [1] J. C. Schuster and J. Bauer, *The Ternary System Titanium-Aluminum-Nitrogen*, J. Solid State Chem. **53**, 260–265 (1984), doi:10.1016/0022-4596(84)90100-2.

## Found in

- [1] K. Cenzual, L. M. Gelato, M. Penzo, and E. Parthé, *Inorganic structure types with revised space groups. I*, Acta Crystallogr. Sect. B **47**, 433–439 (1991), doi:10.1107/S0108768191000903.