


$K_2S_2O_5$ ($K0_1$) Structure: A2B5C2_mP18_11_2e_e2f_2e-001

This structure originally had the label A2B5C2_mP18_11_2e_e2f_2e. Calls to that address will be redirected here.

Cite this page as: D. Hicks, M. J. Mehl, M. Esters, C. Oses, O. Levy, G. L. W. Hart, C. Toher, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 3*, Comput. Mater. Sci. **199**, 110450 (2021), doi: 10.1016/j.commatsci.2021.110450.

<https://aflow.org/p/YYGA>

https://aflow.org/p/A2B5C2_mP18_11_2e_e2f_2e-001




Prototype	$K_2O_5S_2$
AFLOW prototype label	A2B5C2_mP18_11_2e_e2f_2e-001
<i>Strukturbericht</i> designation	$K0_1$
ICSD	60025
Pearson symbol	mP18
Space group number	11
Space group symbol	$P2_1/m$
AFLOW prototype command	<code>aflow --proto=A2B5C2_mP18_11_2e_e2f_2e-001</code> <code>--params=a, b/a, c/a, β, x_1, z_1, x_2, z_2, x_3, z_3, x_4, z_4, x_5, z_5, x_6, y_6, z_6, x_7, y_7, z_7</code>

Other compounds with this structure

$K_2S_2P_5$, $(NH_4)_2S_2P_5$

Simple Monoclinic primitive vectors

$$\begin{aligned}
\mathbf{a}_1 &= a \hat{\mathbf{x}} \\
\mathbf{a}_2 &= b \hat{\mathbf{y}} \\
\mathbf{a}_3 &= c \cos \beta \hat{\mathbf{x}} + c \sin \beta \hat{\mathbf{z}}
\end{aligned}$$



Basis vectors

	Lattice coordinates		Cartesian coordinates	Wyckoff position	Atom type
\mathbf{B}_1	$x_1 \mathbf{a}_1 + \frac{1}{4} \mathbf{a}_2 + z_1 \mathbf{a}_3$	=	$(ax_1 + cz_1 \cos \beta) \hat{\mathbf{x}} + \frac{1}{4}b \hat{\mathbf{y}} + cz_1 \sin \beta \hat{\mathbf{z}}$	(2e)	K I
\mathbf{B}_2	$-x_1 \mathbf{a}_1 + \frac{3}{4} \mathbf{a}_2 - z_1 \mathbf{a}_3$	=	$-(ax_1 + cz_1 \cos \beta) \hat{\mathbf{x}} + \frac{3}{4}b \hat{\mathbf{y}} - cz_1 \sin \beta \hat{\mathbf{z}}$	(2e)	K I
\mathbf{B}_3	$x_2 \mathbf{a}_1 + \frac{1}{4} \mathbf{a}_2 + z_2 \mathbf{a}_3$	=	$(ax_2 + cz_2 \cos \beta) \hat{\mathbf{x}} + \frac{1}{4}b \hat{\mathbf{y}} + cz_2 \sin \beta \hat{\mathbf{z}}$	(2e)	K II
\mathbf{B}_4	$-x_2 \mathbf{a}_1 + \frac{3}{4} \mathbf{a}_2 - z_2 \mathbf{a}_3$	=	$-(ax_2 + cz_2 \cos \beta) \hat{\mathbf{x}} + \frac{3}{4}b \hat{\mathbf{y}} - cz_2 \sin \beta \hat{\mathbf{z}}$	(2e)	K II
\mathbf{B}_5	$x_3 \mathbf{a}_1 + \frac{1}{4} \mathbf{a}_2 + z_3 \mathbf{a}_3$	=	$(ax_3 + cz_3 \cos \beta) \hat{\mathbf{x}} + \frac{1}{4}b \hat{\mathbf{y}} + cz_3 \sin \beta \hat{\mathbf{z}}$	(2e)	O I
\mathbf{B}_6	$-x_3 \mathbf{a}_1 + \frac{3}{4} \mathbf{a}_2 - z_3 \mathbf{a}_3$	=	$-(ax_3 + cz_3 \cos \beta) \hat{\mathbf{x}} + \frac{3}{4}b \hat{\mathbf{y}} - cz_3 \sin \beta \hat{\mathbf{z}}$	(2e)	O I
\mathbf{B}_7	$x_4 \mathbf{a}_1 + \frac{1}{4} \mathbf{a}_2 + z_4 \mathbf{a}_3$	=	$(ax_4 + cz_4 \cos \beta) \hat{\mathbf{x}} + \frac{1}{4}b \hat{\mathbf{y}} + cz_4 \sin \beta \hat{\mathbf{z}}$	(2e)	S I
\mathbf{B}_8	$-x_4 \mathbf{a}_1 + \frac{3}{4} \mathbf{a}_2 - z_4 \mathbf{a}_3$	=	$-(ax_4 + cz_4 \cos \beta) \hat{\mathbf{x}} + \frac{3}{4}b \hat{\mathbf{y}} - cz_4 \sin \beta \hat{\mathbf{z}}$	(2e)	S I
\mathbf{B}_9	$x_5 \mathbf{a}_1 + \frac{1}{4} \mathbf{a}_2 + z_5 \mathbf{a}_3$	=	$(ax_5 + cz_5 \cos \beta) \hat{\mathbf{x}} + \frac{1}{4}b \hat{\mathbf{y}} + cz_5 \sin \beta \hat{\mathbf{z}}$	(2e)	S II
\mathbf{B}_{10}	$-x_5 \mathbf{a}_1 + \frac{3}{4} \mathbf{a}_2 - z_5 \mathbf{a}_3$	=	$-(ax_5 + cz_5 \cos \beta) \hat{\mathbf{x}} + \frac{3}{4}b \hat{\mathbf{y}} - cz_5 \sin \beta \hat{\mathbf{z}}$	(2e)	S II
\mathbf{B}_{11}	$x_6 \mathbf{a}_1 + y_6 \mathbf{a}_2 + z_6 \mathbf{a}_3$	=	$(ax_6 + cz_6 \cos \beta) \hat{\mathbf{x}} + by_6 \hat{\mathbf{y}} + cz_6 \sin \beta \hat{\mathbf{z}}$	(4f)	O II
\mathbf{B}_{12}	$-x_6 \mathbf{a}_1 + (y_6 + \frac{1}{2}) \mathbf{a}_2 - z_6 \mathbf{a}_3$	=	$-(ax_6 + cz_6 \cos \beta) \hat{\mathbf{x}} + b(y_6 + \frac{1}{2}) \hat{\mathbf{y}} - cz_6 \sin \beta \hat{\mathbf{z}}$	(4f)	O II
\mathbf{B}_{13}	$-x_6 \mathbf{a}_1 - y_6 \mathbf{a}_2 - z_6 \mathbf{a}_3$	=	$-(ax_6 + cz_6 \cos \beta) \hat{\mathbf{x}} - by_6 \hat{\mathbf{y}} - cz_6 \sin \beta \hat{\mathbf{z}}$	(4f)	O II
\mathbf{B}_{14}	$x_6 \mathbf{a}_1 - (y_6 - \frac{1}{2}) \mathbf{a}_2 + z_6 \mathbf{a}_3$	=	$(ax_6 + cz_6 \cos \beta) \hat{\mathbf{x}} - b(y_6 - \frac{1}{2}) \hat{\mathbf{y}} + cz_6 \sin \beta \hat{\mathbf{z}}$	(4f)	O II
\mathbf{B}_{15}	$x_7 \mathbf{a}_1 + y_7 \mathbf{a}_2 + z_7 \mathbf{a}_3$	=	$(ax_7 + cz_7 \cos \beta) \hat{\mathbf{x}} + by_7 \hat{\mathbf{y}} + cz_7 \sin \beta \hat{\mathbf{z}}$	(4f)	O III
\mathbf{B}_{16}	$-x_7 \mathbf{a}_1 + (y_7 + \frac{1}{2}) \mathbf{a}_2 - z_7 \mathbf{a}_3$	=	$-(ax_7 + cz_7 \cos \beta) \hat{\mathbf{x}} + b(y_7 + \frac{1}{2}) \hat{\mathbf{y}} - cz_7 \sin \beta \hat{\mathbf{z}}$	(4f)	O III
\mathbf{B}_{17}	$-x_7 \mathbf{a}_1 - y_7 \mathbf{a}_2 - z_7 \mathbf{a}_3$	=	$-(ax_7 + cz_7 \cos \beta) \hat{\mathbf{x}} - by_7 \hat{\mathbf{y}} - cz_7 \sin \beta \hat{\mathbf{z}}$	(4f)	O III
\mathbf{B}_{18}	$x_7 \mathbf{a}_1 - (y_7 - \frac{1}{2}) \mathbf{a}_2 + z_7 \mathbf{a}_3$	=	$(ax_7 + cz_7 \cos \beta) \hat{\mathbf{x}} - b(y_7 - \frac{1}{2}) \hat{\mathbf{y}} + cz_7 \sin \beta \hat{\mathbf{z}}$	(4f)	O III

References

- [1] I.-C. Chen and Y. Wang, *Reinvestigation of potassium pyrosulfite, $K_2S_2O_5$* **40**, 1780–1781 (1984), doi:10.1107/S0108270184009525.