α-RbPr[MoO_4]$_2$ Structure: A2B8CD_oP24_48_h_2m_a_b-001

This structure originally had the label `A2B8CD_oP24_48_k_2m_d_b`. Calls to that address will be redirected here.

Cite this page as: D. Hicks, M. J. Mehl, E. Gossett, C. Toher, O. Levy, R. M. Hanson, G. Hart, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 2*, Comput. Mater. Sci. **161**, S1 (2019). doi: 10.1016/j.commatsci.2018.10.043

https://aflow.org/p/7YK7

https://aflow.org/p/A2B8CD_oP24_48_h_2m_a_b-001



Prototype	Mo_2O_8PrRb
AFLOW prototype label	A2B8CD_oP24_48_h_2m_a_b-001
ICSD	6290
Pearson symbol	oP24
Space group number	48
Space group symbol	*Pnnn*
AFLOW prototype command	`aflow --proto=A2B8CD_oP24_48_h_2m_a_b-001 --params=`$a, b/a, c/a, x_3, x_4, y_4, z_4, x_5, y_5, z_5$

Simple Orthorhombic primitive vectors

$$\mathbf{a_1} = a\,\hat{\mathbf{x}}$$
$$\mathbf{a_2} = b\,\hat{\mathbf{y}}$$
$$\mathbf{a_3} = c\,\hat{\mathbf{z}}$$



Basis vectors

		Lattice coordinates		Cartesian coordinates	Wyckoff position	Atom type
$\mathbf{B_1}$	=	$\frac{1}{4}\,\mathbf{a}_1 + \frac{1}{4}\,\mathbf{a}_2 + \frac{1}{4}\,\mathbf{a}_3$	=	$\frac{1}{4}a\,\hat{\mathbf{x}} + \frac{1}{4}b\,\hat{\mathbf{y}} + \frac{1}{4}c\,\hat{\mathbf{z}}$	(2a)	Pr I
$\mathbf{B_2}$	=	$\frac{3}{4}\,\mathbf{a}_1 + \frac{3}{4}\,\mathbf{a}_2 + \frac{3}{4}\,\mathbf{a}_3$	=	$\frac{3}{4}a\,\hat{\mathbf{x}} + \frac{3}{4}b\,\hat{\mathbf{y}} + \frac{3}{4}c\,\hat{\mathbf{z}}$	(2a)	Pr I
$\mathbf{B_3}$	=	$\frac{3}{4}\,\mathbf{a}_1 + \frac{1}{4}\,\mathbf{a}_2 + \frac{1}{4}\,\mathbf{a}_3$	=	$\frac{3}{4}a\,\hat{\mathbf{x}} + \frac{1}{4}b\,\hat{\mathbf{y}} + \frac{1}{4}c\,\hat{\mathbf{z}}$	(2b)	Rb I
$\mathbf{B_4}$	=	$\frac{1}{4}\,\mathbf{a}_1 + \frac{3}{4}\,\mathbf{a}_2 + \frac{3}{4}\,\mathbf{a}_3$	=	$\frac{1}{4}a\,\hat{\mathbf{x}} + \frac{3}{4}b\,\hat{\mathbf{y}} + \frac{3}{4}c\,\hat{\mathbf{z}}$	(2b)	Rb I
$\mathbf{B_5}$	=	$x_3\,\mathbf{a}_1 + \frac{1}{4}\,\mathbf{a}_2 + \frac{3}{4}\,\mathbf{a}_3$	=	$ax_3\,\hat{\mathbf{x}} + \frac{1}{4}b\,\hat{\mathbf{y}} + \frac{3}{4}c\,\hat{\mathbf{z}}$	(4h)	Mo I
$\mathbf{B_6}$	=	$-\left(x_3 - \frac{1}{2}\right)\,\mathbf{a}_1 + \frac{1}{4}\,\mathbf{a}_2 + \frac{3}{4}\,\mathbf{a}_3$	=	$-a\left(x_3 - \frac{1}{2}\right)\,\hat{\mathbf{x}} + \frac{1}{4}b\,\hat{\mathbf{y}} + \frac{3}{4}c\,\hat{\mathbf{z}}$	(4h)	Mo I
$\mathbf{B_7}$	=	$-x_3\,\mathbf{a}_1 + \frac{3}{4}\,\mathbf{a}_2 + \frac{1}{4}\,\mathbf{a}_3$	=	$-ax_3\,\hat{\mathbf{x}} + \frac{3}{4}b\,\hat{\mathbf{y}} + \frac{1}{4}c\,\hat{\mathbf{z}}$	(4h)	Mo I
$\mathbf{B_8}$	=	$\left(x_3 + \frac{1}{2}\right)\,\mathbf{a}_1 + \frac{3}{4}\,\mathbf{a}_2 + \frac{1}{4}\,\mathbf{a}_3$	=	$a\left(x_3 + \frac{1}{2}\right)\,\hat{\mathbf{x}} + \frac{3}{4}b\,\hat{\mathbf{y}} + \frac{1}{4}c\,\hat{\mathbf{z}}$	(4h)	Mo I
$\mathbf{B_9}$	=	$x_4\,\mathbf{a}_1 + y_4\,\mathbf{a}_2 + z_4\,\mathbf{a}_3$	=	$ax_4\,\hat{\mathbf{x}} + by_4\,\hat{\mathbf{y}} + cz_4\,\hat{\mathbf{z}}$	(8m)	O I
$\mathbf{B_{10}}$	=	$-\left(x_4 - \frac{1}{2}\right)\,\mathbf{a}_1 - \left(y_4 - \frac{1}{2}\right)\,\mathbf{a}_2 + z_4\,\mathbf{a}_3$	=	$-a\left(x_4 - \frac{1}{2}\right)\,\hat{\mathbf{x}} - b\left(y_4 - \frac{1}{2}\right)\,\hat{\mathbf{y}} + cz_4\,\hat{\mathbf{z}}$	(8m)	O I
$\mathbf{B_{11}}$	=	$-\left(x_4 - \frac{1}{2}\right)\,\mathbf{a}_1 + y_4\,\mathbf{a}_2 - \left(z_4 - \frac{1}{2}\right)\,\mathbf{a}_3$	=	$-a\left(x_4 - \frac{1}{2}\right)\,\hat{\mathbf{x}} + by_4\,\hat{\mathbf{y}} - c\left(z_4 - \frac{1}{2}\right)\,\hat{\mathbf{z}}$	(8m)	O I
$\mathbf{B_{12}}$	=	$x_4\,\mathbf{a}_1 - \left(y_4 - \frac{1}{2}\right)\,\mathbf{a}_2 - \left(z_4 - \frac{1}{2}\right)\,\mathbf{a}_3$	=	$ax_4\,\hat{\mathbf{x}} - b\left(y_4 - \frac{1}{2}\right)\,\hat{\mathbf{y}} - c\left(z_4 - \frac{1}{2}\right)\,\hat{\mathbf{z}}$	(8m)	O I
$\mathbf{B_{13}}$	=	$-x_4\,\mathbf{a}_1 - y_4\,\mathbf{a}_2 - z_4\,\mathbf{a}_3$	=	$-ax_4\,\hat{\mathbf{x}} - by_4\,\hat{\mathbf{y}} - cz_4\,\hat{\mathbf{z}}$	(8m)	O I
$\mathbf{B_{14}}$	=	$\left(x_4 + \frac{1}{2}\right)\,\mathbf{a}_1 + \left(y_4 + \frac{1}{2}\right)\,\mathbf{a}_2 - z_4\,\mathbf{a}_3$	=	$a\left(x_4 + \frac{1}{2}\right)\,\hat{\mathbf{x}} + b\left(y_4 + \frac{1}{2}\right)\,\hat{\mathbf{y}} - cz_4\,\hat{\mathbf{z}}$	(8m)	O I
$\mathbf{B_{15}}$	=	$\left(x_4 + \frac{1}{2}\right)\,\mathbf{a}_1 - y_4\,\mathbf{a}_2 + \left(z_4 + \frac{1}{2}\right)\,\mathbf{a}_3$	=	$a\left(x_4 + \frac{1}{2}\right)\,\hat{\mathbf{x}} - by_4\,\hat{\mathbf{y}} + c\left(z_4 + \frac{1}{2}\right)\,\hat{\mathbf{z}}$	(8m)	O I
$\mathbf{B_{16}}$	=	$-x_4\,\mathbf{a}_1 + \left(y_4 + \frac{1}{2}\right)\,\mathbf{a}_2 + \left(z_4 + \frac{1}{2}\right)\,\mathbf{a}_3$	=	$-ax_4\,\hat{\mathbf{x}} + b\left(y_4 + \frac{1}{2}\right)\,\hat{\mathbf{y}} + c\left(z_4 + \frac{1}{2}\right)\,\hat{\mathbf{z}}$	(8m)	O I
$\mathbf{B_{17}}$	=	$x_5\,\mathbf{a}_1 + y_5\,\mathbf{a}_2 + z_5\,\mathbf{a}_3$	=	$ax_5\,\hat{\mathbf{x}} + by_5\,\hat{\mathbf{y}} + cz_5\,\hat{\mathbf{z}}$	(8m)	O II
$\mathbf{B_{18}}$	=	$-\left(x_5 - \frac{1}{2}\right)\,\mathbf{a}_1 - \left(y_5 - \frac{1}{2}\right)\,\mathbf{a}_2 + z_5\,\mathbf{a}_3$	=	$-a\left(x_5 - \frac{1}{2}\right)\,\hat{\mathbf{x}} - b\left(y_5 - \frac{1}{2}\right)\,\hat{\mathbf{y}} + cz_5\,\hat{\mathbf{z}}$	(8m)	O II
$\mathbf{B_{19}}$	=	$-\left(x_5 - \frac{1}{2}\right)\,\mathbf{a}_1 + y_5\,\mathbf{a}_2 - \left(z_5 - \frac{1}{2}\right)\,\mathbf{a}_3$	=	$-a\left(x_5 - \frac{1}{2}\right)\,\hat{\mathbf{x}} + by_5\,\hat{\mathbf{y}} - c\left(z_5 - \frac{1}{2}\right)\,\hat{\mathbf{z}}$	(8m)	O II
$\mathbf{B_{20}}$	=	$x_5\,\mathbf{a}_1 - \left(y_5 - \frac{1}{2}\right)\,\mathbf{a}_2 - \left(z_5 - \frac{1}{2}\right)\,\mathbf{a}_3$	=	$ax_5\,\hat{\mathbf{x}} - b\left(y_5 - \frac{1}{2}\right)\,\hat{\mathbf{y}} - c\left(z_5 - \frac{1}{2}\right)\,\hat{\mathbf{z}}$	(8m)	O II
$\mathbf{B_{21}}$	=	$-x_5\,\mathbf{a}_1 - y_5\,\mathbf{a}_2 - z_5\,\mathbf{a}_3$	=	$-ax_5\,\hat{\mathbf{x}} - by_5\,\hat{\mathbf{y}} - cz_5\,\hat{\mathbf{z}}$	(8m)	O II
$\mathbf{B_{22}}$	=	$\left(x_5 + \frac{1}{2}\right)\,\mathbf{a}_1 + \left(y_5 + \frac{1}{2}\right)\,\mathbf{a}_2 - z_5\,\mathbf{a}_3$	=	$a\left(x_5 + \frac{1}{2}\right)\,\hat{\mathbf{x}} + b\left(y_5 + \frac{1}{2}\right)\,\hat{\mathbf{y}} - cz_5\,\hat{\mathbf{z}}$	(8m)	O II
$\mathbf{B_{23}}$	=	$\left(x_5 + \frac{1}{2}\right)\,\mathbf{a}_1 - y_5\,\mathbf{a}_2 + \left(z_5 + \frac{1}{2}\right)\,\mathbf{a}_3$	=	$a\left(x_5 + \frac{1}{2}\right)\,\hat{\mathbf{x}} - by_5\,\hat{\mathbf{y}} + c\left(z_5 + \frac{1}{2}\right)\,\hat{\mathbf{z}}$	(8m)	O II
$\mathbf{B_{24}}$	=	$-x_5\,\mathbf{a}_1 + \left(y_5 + \frac{1}{2}\right)\,\mathbf{a}_2 + \left(z_5 + \frac{1}{2}\right)\,\mathbf{a}_3$	=	$-ax_5\,\hat{\mathbf{x}} + b\left(y_5 + \frac{1}{2}\right)\,\hat{\mathbf{y}} + c\left(z_5 + \frac{1}{2}\right)\,\hat{\mathbf{z}}$	(8m)	O II

References

[1] R. F. Klevtsova and P. V. Klevtsov, *Polymorphism of rubidium–praseodymium molybdate, $RbPr(MoO_4)_2$*, Kristallografiya **15**, 466–470 (1970).

Found in

[1] P. Villars and K. Cenzual, *Pearson's Crystal Data – Crystal Structure Database for Inorganic Compounds* (2013). ASM International.