


β -RuCl₃ Structure: A3B_hP8_185_c_a-001

This structure originally had the label **A3B_hP8_185_c_a**. Calls to that address will be redirected here.

Cite this page as: D. Hicks, M. J. Mehl, E. Gossett, C. Toher, O. Levy, R. M. Hanson, G. Hart, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 2*, Comput. Mater. Sci. **161**, S1 (2019). doi: 10.1016/j.commatsci.2018.10.043

<https://aflow.org/p/5RVU>

https://aflow.org/p/A3B_hP8_185_c_a-001




Prototype	Cl ₃ Ru
AFLOW prototype label	A3B_hP8_185_c_a-001
ICSD	22092
Pearson symbol	hP8
Space group number	185
Space group symbol	$P6_3cm$
AFLOW prototype command	<code>aflow --proto=A3B_hP8_185_c_a-001 --params=a, c/a, z₁, x₂, z₂</code>

- The crystal structure of both α - and β - RuCl_3 is somewhat uncertain:
- α - RuCl_3 has been reported in the trigonal CrCl_3 ($D0_4$) structure (Fletcher, 1967) and in the monoclinic AlCl_3 structure (Johnson, 2015).
- (Fletcher, 1967) states that the structure of β - RuCl_3 is consistent with the hexagonal space groups $P6_3cm$ #185 (this structure), $P6c2$ #188, and $P6_3/mcm$ #193, in addition to the trigonal space group $P3c1$ #158, which we present here: trigonal β - RuCl_3 (A3B.hp8_158_d.a).
- Space group $P6_3cm$ #185 allows an arbitrary choice of the origin of the z -axis. Here we set $z_1 = 0$ for the ruthenium (2a) atoms.

Hexagonal primitive vectors

$$\begin{aligned}\mathbf{a}_1 &= \frac{1}{2}a \hat{\mathbf{x}} - \frac{\sqrt{3}}{2}a \hat{\mathbf{y}} \\ \mathbf{a}_2 &= \frac{1}{2}a \hat{\mathbf{x}} + \frac{\sqrt{3}}{2}a \hat{\mathbf{y}} \\ \mathbf{a}_3 &= c \hat{\mathbf{z}}\end{aligned}$$



Basis vectors

	Lattice coordinates		Cartesian coordinates	Wyckoff position	Atom type
\mathbf{B}_1	$z_1 \mathbf{a}_3$	=	$cz_1 \hat{\mathbf{z}}$	(2a)	Ru I
\mathbf{B}_2	$(z_1 + \frac{1}{2}) \mathbf{a}_3$	=	$c(z_1 + \frac{1}{2}) \hat{\mathbf{z}}$	(2a)	Ru I
\mathbf{B}_3	$x_2 \mathbf{a}_1 + z_2 \mathbf{a}_3$	=	$\frac{1}{2}ax_2 \hat{\mathbf{x}} - \frac{\sqrt{3}}{2}ax_2 \hat{\mathbf{y}} + cz_2 \hat{\mathbf{z}}$	(6c)	Cl I
\mathbf{B}_4	$x_2 \mathbf{a}_2 + z_2 \mathbf{a}_3$	=	$\frac{1}{2}ax_2 \hat{\mathbf{x}} + \frac{\sqrt{3}}{2}ax_2 \hat{\mathbf{y}} + cz_2 \hat{\mathbf{z}}$	(6c)	Cl I
\mathbf{B}_5	$-x_2 \mathbf{a}_1 - x_2 \mathbf{a}_2 + z_2 \mathbf{a}_3$	=	$-ax_2 \hat{\mathbf{x}} + cz_2 \hat{\mathbf{z}}$	(6c)	Cl I
\mathbf{B}_6	$-x_2 \mathbf{a}_1 + (z_2 + \frac{1}{2}) \mathbf{a}_3$	=	$-\frac{1}{2}ax_2 \hat{\mathbf{x}} + \frac{\sqrt{3}}{2}ax_2 \hat{\mathbf{y}} + c(z_2 + \frac{1}{2}) \hat{\mathbf{z}}$	(6c)	Cl I
\mathbf{B}_7	$-x_2 \mathbf{a}_2 + (z_2 + \frac{1}{2}) \mathbf{a}_3$	=	$-\frac{1}{2}ax_2 \hat{\mathbf{x}} - \frac{\sqrt{3}}{2}ax_2 \hat{\mathbf{y}} + c(z_2 + \frac{1}{2}) \hat{\mathbf{z}}$	(6c)	Cl I
\mathbf{B}_8	$x_2 \mathbf{a}_1 + x_2 \mathbf{a}_2 + (z_2 + \frac{1}{2}) \mathbf{a}_3$	=	$ax_2 \hat{\mathbf{x}} + c(z_2 + \frac{1}{2}) \hat{\mathbf{z}}$	(6c)	Cl I

References

- [1] J. M. Fletcher, W. E. Gardner, A. C. Fox, and G. Topping, *X-Ray, infrared, and magnetic studies of α - and β -ruthenium trichloride*, J. Chem. Soc. A pp. 1038–1045 (1967), doi:10.1039/J19670001038.