Rh_2Ga_9 Structure: A9B2_mP22_7_9a_2a-001

This structure originally had the label `A9B2_mP22_7_9a_2a`. Calls to that address will be redirected here.

Cite this page as: D. Hicks, M. J. Mehl, E. Gossett, C. Toher, O. Levy, R. M. Hanson, G. Hart, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 2*, Comput. Mater. Sci. **161**, S1 (2019). doi: 10.1016/j.commatsci.2018.10.043

https://aflow.org/p/2DRY

https://aflow.org/p/A9B2_mP22_7_9a_2a-001



Prototype	Ga_9Rh_2
AFLOW prototype label	A9B2_mP22_7_9a_2a-001
ICSD	414305
Pearson symbol	mP22
Space group number	7
Space group symbol	Pc
AFLOW prototype command	`aflow --proto=A9B2_mP22_7_9a_2a-001 --params=`$a, b/a, c/a, \beta, x_1, y_1, z_1, x_2, y_2, z_2, x_3, y_3, z_3, x_4, y_4, z_4, x_5, y_5, z_5, x_6, y_6, z_6, x_7, y_7, z_7, x_8, y_8, z_8, x_9, y_9, z_9, x_{10}, y_{10}, z_{10}, x_{11}, y_{11}, z_{11}$

Other compounds with this structure

Ir_2Ga_9

Simple Monoclinic primitive vectors

$$
\begin{aligned}
\mathbf{a_1} &= a\,\hat{\mathbf{x}} \\
\mathbf{a_2} &= b\,\hat{\mathbf{y}} \\
\mathbf{a_3} &= c\cos\beta\,\hat{\mathbf{x}} + c\sin\beta\,\hat{\mathbf{z}}
\end{aligned}
$$



Basis vectors

		Lattice coordinates		Cartesian coordinates	Wyckoff position	Atom type
$\mathbf{B_1}$	=	$x_1\,\mathbf{a}_1 + y_1\,\mathbf{a}_2 + z_1\,\mathbf{a}_3$	=	$(ax_1 + cz_1\cos\beta)\,\hat{\mathbf{x}} + by_1\,\hat{\mathbf{y}} + cz_1\sin\beta\,\hat{\mathbf{z}}$	(2a)	Ga I
$\mathbf{B_2}$	=	$x_1\,\mathbf{a}_1 - y_1\,\mathbf{a}_2 + \left(z_1 + \frac{1}{2}\right)\,\mathbf{a}_3$	=	$\left(ax_1 + c\left(z_1 + \frac{1}{2}\right)\cos\beta\right)\,\hat{\mathbf{x}} - by_1\,\hat{\mathbf{y}} + c\left(z_1 + \frac{1}{2}\right)\sin\beta\,\hat{\mathbf{z}}$	(2a)	Ga I
$\mathbf{B_3}$	=	$x_2\,\mathbf{a}_1 + y_2\,\mathbf{a}_2 + z_2\,\mathbf{a}_3$	=	$(ax_2 + cz_2\cos\beta)\,\hat{\mathbf{x}} + by_2\,\hat{\mathbf{y}} + cz_2\sin\beta\,\hat{\mathbf{z}}$	(2a)	Ga II
$\mathbf{B_4}$	=	$x_2\,\mathbf{a}_1 - y_2\,\mathbf{a}_2 + \left(z_2 + \frac{1}{2}\right)\,\mathbf{a}_3$	=	$\left(ax_2 + c\left(z_2 + \frac{1}{2}\right)\cos\beta\right)\,\hat{\mathbf{x}} - by_2\,\hat{\mathbf{y}} + c\left(z_2 + \frac{1}{2}\right)\sin\beta\,\hat{\mathbf{z}}$	(2a)	Ga II
$\mathbf{B_5}$	=	$x_3\,\mathbf{a}_1 + y_3\,\mathbf{a}_2 + z_3\,\mathbf{a}_3$	=	$(ax_3 + cz_3\cos\beta)\,\hat{\mathbf{x}} + by_3\,\hat{\mathbf{y}} + cz_3\sin\beta\,\hat{\mathbf{z}}$	(2a)	Ga III
$\mathbf{B_6}$	=	$x_3\,\mathbf{a}_1 - y_3\,\mathbf{a}_2 + \left(z_3 + \frac{1}{2}\right)\,\mathbf{a}_3$	=	$\left(ax_3 + c\left(z_3 + \frac{1}{2}\right)\cos\beta\right)\,\hat{\mathbf{x}} - by_3\,\hat{\mathbf{y}} + c\left(z_3 + \frac{1}{2}\right)\sin\beta\,\hat{\mathbf{z}}$	(2a)	Ga III
$\mathbf{B_7}$	=	$x_4\,\mathbf{a}_1 + y_4\,\mathbf{a}_2 + z_4\,\mathbf{a}_3$	=	$(ax_4 + cz_4\cos\beta)\,\hat{\mathbf{x}} + by_4\,\hat{\mathbf{y}} + cz_4\sin\beta\,\hat{\mathbf{z}}$	(2a)	Ga IV
$\mathbf{B_8}$	=	$x_4\,\mathbf{a}_1 - y_4\,\mathbf{a}_2 + \left(z_4 + \frac{1}{2}\right)\,\mathbf{a}_3$	=	$\left(ax_4 + c\left(z_4 + \frac{1}{2}\right)\cos\beta\right)\,\hat{\mathbf{x}} - by_4\,\hat{\mathbf{y}} + c\left(z_4 + \frac{1}{2}\right)\sin\beta\,\hat{\mathbf{z}}$	(2a)	Ga IV
$\mathbf{B_9}$	=	$x_5\,\mathbf{a}_1 + y_5\,\mathbf{a}_2 + z_5\,\mathbf{a}_3$	=	$(ax_5 + cz_5\cos\beta)\,\hat{\mathbf{x}} + by_5\,\hat{\mathbf{y}} + cz_5\sin\beta\,\hat{\mathbf{z}}$	(2a)	Ga V
$\mathbf{B_{10}}$	=	$x_5\,\mathbf{a}_1 - y_5\,\mathbf{a}_2 + \left(z_5 + \frac{1}{2}\right)\,\mathbf{a}_3$	=	$\left(ax_5 + c\left(z_5 + \frac{1}{2}\right)\cos\beta\right)\,\hat{\mathbf{x}} - by_5\,\hat{\mathbf{y}} + c\left(z_5 + \frac{1}{2}\right)\sin\beta\,\hat{\mathbf{z}}$	(2a)	Ga V
$\mathbf{B_{11}}$	=	$x_6\,\mathbf{a}_1 + y_6\,\mathbf{a}_2 + z_6\,\mathbf{a}_3$	=	$(ax_6 + cz_6\cos\beta)\,\hat{\mathbf{x}} + by_6\,\hat{\mathbf{y}} + cz_6\sin\beta\,\hat{\mathbf{z}}$	(2a)	Ga VI
$\mathbf{B_{12}}$	=	$x_6\,\mathbf{a}_1 - y_6\,\mathbf{a}_2 + \left(z_6 + \frac{1}{2}\right)\,\mathbf{a}_3$	=	$\left(ax_6 + c\left(z_6 + \frac{1}{2}\right)\cos\beta\right)\,\hat{\mathbf{x}} - by_6\,\hat{\mathbf{y}} + c\left(z_6 + \frac{1}{2}\right)\sin\beta\,\hat{\mathbf{z}}$	(2a)	Ga VI
$\mathbf{B_{13}}$	=	$x_7\,\mathbf{a}_1 + y_7\,\mathbf{a}_2 + z_7\,\mathbf{a}_3$	=	$(ax_7 + cz_7\cos\beta)\,\hat{\mathbf{x}} + by_7\,\hat{\mathbf{y}} + cz_7\sin\beta\,\hat{\mathbf{z}}$	(2a)	Ga VII
$\mathbf{B_{14}}$	=	$x_7\,\mathbf{a}_1 - y_7\,\mathbf{a}_2 + \left(z_7 + \frac{1}{2}\right)\,\mathbf{a}_3$	=	$\left(ax_7 + c\left(z_7 + \frac{1}{2}\right)\cos\beta\right)\,\hat{\mathbf{x}} - by_7\,\hat{\mathbf{y}} + c\left(z_7 + \frac{1}{2}\right)\sin\beta\,\hat{\mathbf{z}}$	(2a)	Ga VII
$\mathbf{B_{15}}$	=	$x_8\,\mathbf{a}_1 + y_8\,\mathbf{a}_2 + z_8\,\mathbf{a}_3$	=	$(ax_8 + cz_8\cos\beta)\,\hat{\mathbf{x}} + by_8\,\hat{\mathbf{y}} + cz_8\sin\beta\,\hat{\mathbf{z}}$	(2a)	Ga VIII
$\mathbf{B_{16}}$	=	$x_8\,\mathbf{a}_1 - y_8\,\mathbf{a}_2 + \left(z_8 + \frac{1}{2}\right)\,\mathbf{a}_3$	=	$\left(ax_8 + c\left(z_8 + \frac{1}{2}\right)\cos\beta\right)\,\hat{\mathbf{x}} - by_8\,\hat{\mathbf{y}} + c\left(z_8 + \frac{1}{2}\right)\sin\beta\,\hat{\mathbf{z}}$	(2a)	Ga VIII
$\mathbf{B_{17}}$	=	$x_9\,\mathbf{a}_1 + y_9\,\mathbf{a}_2 + z_9\,\mathbf{a}_3$	=	$(ax_9 + cz_9\cos\beta)\,\hat{\mathbf{x}} + by_9\,\hat{\mathbf{y}} + cz_9\sin\beta\,\hat{\mathbf{z}}$	(2a)	Ga IX
$\mathbf{B_{18}}$	=	$x_9\,\mathbf{a}_1 - y_9\,\mathbf{a}_2 + \left(z_9 + \frac{1}{2}\right)\,\mathbf{a}_3$	=	$\left(ax_9 + c\left(z_9 + \frac{1}{2}\right)\cos\beta\right)\,\hat{\mathbf{x}} - by_9\,\hat{\mathbf{y}} + c\left(z_9 + \frac{1}{2}\right)\sin\beta\,\hat{\mathbf{z}}$	(2a)	Ga IX
$\mathbf{B_{19}}$	=	$x_{10}\,\mathbf{a}_1 + y_{10}\,\mathbf{a}_2 + z_{10}\,\mathbf{a}_3$	=	$(ax_{10} + cz_{10}\cos\beta)\,\hat{\mathbf{x}} + by_{10}\,\hat{\mathbf{y}} + cz_{10}\sin\beta\,\hat{\mathbf{z}}$	(2a)	Rh I
$\mathbf{B_{20}}$	=	$x_{10}\,\mathbf{a}_1 - y_{10}\,\mathbf{a}_2 + \left(z_{10} + \frac{1}{2}\right)\,\mathbf{a}_3$	=	$\left(ax_{10} + c\left(z_{10} + \frac{1}{2}\right)\cos\beta\right)\,\hat{\mathbf{x}} - by_{10}\,\hat{\mathbf{y}} + c\left(z_{10} + \frac{1}{2}\right)\sin\beta\,\hat{\mathbf{z}}$	(2a)	Rh I
$\mathbf{B_{21}}$	=	$x_{11}\,\mathbf{a}_1 + y_{11}\,\mathbf{a}_2 + z_{11}\,\mathbf{a}_3$	=	$(ax_{11} + cz_{11}\cos\beta)\,\hat{\mathbf{x}} + by_{11}\,\hat{\mathbf{y}} + cz_{11}\sin\beta\,\hat{\mathbf{z}}$	(2a)	Rh II
$\mathbf{B_{22}}$	=	$x_{11}\,\mathbf{a}_1 - y_{11}\,\mathbf{a}_2 + \left(z_{11} + \frac{1}{2}\right)\,\mathbf{a}_3$	=	$\left(ax_{11} + c\left(z_{11} + \frac{1}{2}\right)\cos\beta\right)\,\hat{\mathbf{x}} - by_{11}\,\hat{\mathbf{y}} + c\left(z_{11} + \frac{1}{2}\right)\sin\beta\,\hat{\mathbf{z}}$	(2a)	Rh II

References

[1] M. Boström, H. Rosner, Y. Prots, U. Burkhardt, and Y. Grin, *The Co_2Al_9 Structure Type Revisited*, Z. Anorganische und Allgemeine Chemie **631**, 534–541 (2005), doi:10.1002/zaac.200400418.

Found in

[1] P. Villars and K. Cenzual, *Pearson's Crystal Data – Crystal Structure Database for Inorganic Compounds* (2013). ASM International.