


Chalcanthite ($\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, $H4_{10}$) Structure: AB10C9D_aP42_2_bc_10i_9i_i-001

This structure originally had the label AB10C9D_aP42_2_ae_10i_9i_i. Calls to that address will be redirected here.

Cite this page as: D. Hicks, M. J. Mehl, M. Esters, C. Oses, O. Levy, G. L. W. Hart, C. Toher, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 3*, Comput. Mater. Sci. **199**, 110450 (2021), doi: 10.1016/j.commatsci.2021.110450.

<https://aflow.org/p/NQQF>

https://aflow.org/p/AB10C9D_aP42_2_bc_10i_9i_i-001




Prototype	$\text{CuH}_{10}\text{O}_9\text{S}$
AFLOW prototype label	AB10C9D_aP42_2_bc_10i_9i_i-001
<i>Strukturbericht</i> designation	$H4_{10}$
Mineral name	chalcanthite
ICSD	4305
Pearson symbol	aP42
Space group number	2
Space group symbol	$P\bar{1}$
AFLOW prototype command	<pre>aflow --proto=AB10C9D_aP42_2_bc_10i_9i_i-001 --params=a, b/a, c/a, α, β, γ, $x_3, y_3, z_3, x_4, y_4, z_4, x_5, y_5, z_5, x_6, y_6, z_6, x_7, y_7, z_7, x_8, y_8, z_8, x_9, y_9, z_9, x_{10}, y_{10}, z_{10}, x_{11}, y_{11}, z_{11}, x_{12}, y_{12}, z_{12}, x_{13}, y_{13}, z_{13}, x_{14}, y_{14}, z_{14}, x_{15}, y_{15}, z_{15}, x_{16}, y_{16}, z_{16}, x_{17}, y_{17}, z_{17}, x_{18}, y_{18}, z_{18}, x_{19}, y_{19}, z_{19}, x_{20}, y_{20}, z_{20}, x_{21}, y_{21}, z_{21}, x_{22}, y_{22}, z_{22}$</pre>

- We reference (Bacon, 1975) in order to include the positions of the hydrogen atoms.

Triclinic primitive vectors

$$\begin{aligned}
\mathbf{a}_1 &= a \hat{\mathbf{x}} \\
\mathbf{a}_2 &= b \cos \gamma \hat{\mathbf{x}} + b \sin \gamma \hat{\mathbf{y}} \\
\mathbf{a}_3 &= c_x \hat{\mathbf{x}} + c_y \hat{\mathbf{y}} + c_z \hat{\mathbf{z}} \\
c_x &= c \cos \beta \\
c_y &= c(\cos \alpha - \cos \beta \cos \gamma) / \sin \gamma \\
c_z &= \sqrt{c^2 - c_x^2 - c_y^2}
\end{aligned}$$



Basis vectors

	Lattice coordinates		Cartesian coordinates	Wyckoff position	Atom type
\mathbf{B}_1	$= \frac{1}{2} \mathbf{a}_3$	$=$	$\frac{1}{2} c_x \hat{\mathbf{x}} + \frac{1}{2} c_y \hat{\mathbf{y}} + \frac{1}{2} c_z \hat{\mathbf{z}}$	(1b)	Cu I
\mathbf{B}_2	$= \frac{1}{2} \mathbf{a}_2$	$=$	$\frac{1}{2} b \cos \gamma \hat{\mathbf{x}} + \frac{1}{2} b \sin \gamma \hat{\mathbf{y}}$	(1c)	Cu II
\mathbf{B}_3	$= x_3 \mathbf{a}_1 + y_3 \mathbf{a}_2 + z_3 \mathbf{a}_3$	$=$	$(ax_3 + by_3 \cos \gamma + c_x z_3) \hat{\mathbf{x}} + (by_3 \sin \gamma + c_y z_3) \hat{\mathbf{y}} + c_z z_3 \hat{\mathbf{z}}$	(2i)	H I
\mathbf{B}_4	$= -x_3 \mathbf{a}_1 - y_3 \mathbf{a}_2 - z_3 \mathbf{a}_3$	$=$	$-(ax_3 + by_3 \cos \gamma + c_x z_3) \hat{\mathbf{x}} - (by_3 \sin \gamma + c_y z_3) \hat{\mathbf{y}} - c_z z_3 \hat{\mathbf{z}}$	(2i)	H I
\mathbf{B}_5	$= x_4 \mathbf{a}_1 + y_4 \mathbf{a}_2 + z_4 \mathbf{a}_3$	$=$	$(ax_4 + by_4 \cos \gamma + c_x z_4) \hat{\mathbf{x}} + (by_4 \sin \gamma + c_y z_4) \hat{\mathbf{y}} + c_z z_4 \hat{\mathbf{z}}$	(2i)	H II
\mathbf{B}_6	$= -x_4 \mathbf{a}_1 - y_4 \mathbf{a}_2 - z_4 \mathbf{a}_3$	$=$	$-(ax_4 + by_4 \cos \gamma + c_x z_4) \hat{\mathbf{x}} - (by_4 \sin \gamma + c_y z_4) \hat{\mathbf{y}} - c_z z_4 \hat{\mathbf{z}}$	(2i)	H II
\mathbf{B}_7	$= x_5 \mathbf{a}_1 + y_5 \mathbf{a}_2 + z_5 \mathbf{a}_3$	$=$	$(ax_5 + by_5 \cos \gamma + c_x z_5) \hat{\mathbf{x}} + (by_5 \sin \gamma + c_y z_5) \hat{\mathbf{y}} + c_z z_5 \hat{\mathbf{z}}$	(2i)	H III
\mathbf{B}_8	$= -x_5 \mathbf{a}_1 - y_5 \mathbf{a}_2 - z_5 \mathbf{a}_3$	$=$	$-(ax_5 + by_5 \cos \gamma + c_x z_5) \hat{\mathbf{x}} - (by_5 \sin \gamma + c_y z_5) \hat{\mathbf{y}} - c_z z_5 \hat{\mathbf{z}}$	(2i)	H III
\mathbf{B}_9	$= x_6 \mathbf{a}_1 + y_6 \mathbf{a}_2 + z_6 \mathbf{a}_3$	$=$	$(ax_6 + by_6 \cos \gamma + c_x z_6) \hat{\mathbf{x}} + (by_6 \sin \gamma + c_y z_6) \hat{\mathbf{y}} + c_z z_6 \hat{\mathbf{z}}$	(2i)	H IV
\mathbf{B}_{10}	$= -x_6 \mathbf{a}_1 - y_6 \mathbf{a}_2 - z_6 \mathbf{a}_3$	$=$	$-(ax_6 + by_6 \cos \gamma + c_x z_6) \hat{\mathbf{x}} - (by_6 \sin \gamma + c_y z_6) \hat{\mathbf{y}} - c_z z_6 \hat{\mathbf{z}}$	(2i)	H IV
\mathbf{B}_{11}	$= x_7 \mathbf{a}_1 + y_7 \mathbf{a}_2 + z_7 \mathbf{a}_3$	$=$	$(ax_7 + by_7 \cos \gamma + c_x z_7) \hat{\mathbf{x}} + (by_7 \sin \gamma + c_y z_7) \hat{\mathbf{y}} + c_z z_7 \hat{\mathbf{z}}$	(2i)	H V
\mathbf{B}_{12}	$= -x_7 \mathbf{a}_1 - y_7 \mathbf{a}_2 - z_7 \mathbf{a}_3$	$=$	$-(ax_7 + by_7 \cos \gamma + c_x z_7) \hat{\mathbf{x}} - (by_7 \sin \gamma + c_y z_7) \hat{\mathbf{y}} - c_z z_7 \hat{\mathbf{z}}$	(2i)	H V
\mathbf{B}_{13}	$= x_8 \mathbf{a}_1 + y_8 \mathbf{a}_2 + z_8 \mathbf{a}_3$	$=$	$(ax_8 + by_8 \cos \gamma + c_x z_8) \hat{\mathbf{x}} + (by_8 \sin \gamma + c_y z_8) \hat{\mathbf{y}} + c_z z_8 \hat{\mathbf{z}}$	(2i)	H VI
\mathbf{B}_{14}	$= -x_8 \mathbf{a}_1 - y_8 \mathbf{a}_2 - z_8 \mathbf{a}_3$	$=$	$-(ax_8 + by_8 \cos \gamma + c_x z_8) \hat{\mathbf{x}} - (by_8 \sin \gamma + c_y z_8) \hat{\mathbf{y}} - c_z z_8 \hat{\mathbf{z}}$	(2i)	H VI
\mathbf{B}_{15}	$= x_9 \mathbf{a}_1 + y_9 \mathbf{a}_2 + z_9 \mathbf{a}_3$	$=$	$(ax_9 + by_9 \cos \gamma + c_x z_9) \hat{\mathbf{x}} + (by_9 \sin \gamma + c_y z_9) \hat{\mathbf{y}} + c_z z_9 \hat{\mathbf{z}}$	(2i)	H VII
\mathbf{B}_{16}	$= -x_9 \mathbf{a}_1 - y_9 \mathbf{a}_2 - z_9 \mathbf{a}_3$	$=$	$-(ax_9 + by_9 \cos \gamma + c_x z_9) \hat{\mathbf{x}} - (by_9 \sin \gamma + c_y z_9) \hat{\mathbf{y}} - c_z z_9 \hat{\mathbf{z}}$	(2i)	H VII
\mathbf{B}_{17}	$= x_{10} \mathbf{a}_1 + y_{10} \mathbf{a}_2 + z_{10} \mathbf{a}_3$	$=$	$(ax_{10} + by_{10} \cos \gamma + c_x z_{10}) \hat{\mathbf{x}} + (by_{10} \sin \gamma + c_y z_{10}) \hat{\mathbf{y}} + c_z z_{10} \hat{\mathbf{z}}$	(2i)	H VIII
\mathbf{B}_{18}	$= -x_{10} \mathbf{a}_1 - y_{10} \mathbf{a}_2 - z_{10} \mathbf{a}_3$	$=$	$-(ax_{10} + by_{10} \cos \gamma + c_x z_{10}) \hat{\mathbf{x}} - (by_{10} \sin \gamma + c_y z_{10}) \hat{\mathbf{y}} - c_z z_{10} \hat{\mathbf{z}}$	(2i)	H VIII
\mathbf{B}_{19}	$= x_{11} \mathbf{a}_1 + y_{11} \mathbf{a}_2 + z_{11} \mathbf{a}_3$	$=$	$(ax_{11} + by_{11} \cos \gamma + c_x z_{11}) \hat{\mathbf{x}} + (by_{11} \sin \gamma + c_y z_{11}) \hat{\mathbf{y}} + c_z z_{11} \hat{\mathbf{z}}$	(2i)	H IX

References

- [1] G. E. Bacon and D. H. Titterton, *Neutron-diffraction studies of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ and $\text{CuSO}_4 \cdot 5\text{D}_2\text{O}$* , *Z. Kristallogr.* **141**, 330–341 (1975), doi:10.1524/zkri.1975.141.16.330.

Found in

- [1] R. T. Downs and M. Hall-Wallace, *The American Mineralogist Crystal Structure Database*, *Am. Mineral.* **88**, 247–250 (2003).