Miargyrite ($AgSbS_2$) Structure: AB2C_mC32_15_ae_2f_f-001

Cite this page as: H. Eckert, S. Divilov, A. Zettel, M. J. Mehl, D. Hicks, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 4*. In preparation.

https://aflow.org/p/YUVG

https://aflow.org/p/AB2C_mC32_15_ae_2f_f-001



Prototype	AgS_2Sb
AFLOW prototype label	AB2C_mC32_15_ae_2f_f-001
Mineral name	miargyrite
ICSD	94647
Pearson symbol	mC32
Space group number	15
Space group symbol	$C2/c$
AFLOW prototype command	`aflow --proto=AB2C_mC32_15_ae_2f_f-001 --params=`$a, b/a, c/a, \beta, y_2, x_3, y_3, z_3, x_4, y_4, z_4, x_5, y_5, z_5$

- We shifted the origin of the structure described by (Effenberger, 2002) so that a silver atom is at the origin, the Wyckoff (4a) site, rather than at (0 1/2 1/2), Wyckoff (4b).

Base-centered Monoclinic primitive vectors

$$\mathbf{a_1} = \frac{1}{2}a\,\hat{\mathbf{x}} - \frac{1}{2}b\,\hat{\mathbf{y}}$$
$$\mathbf{a_2} = \frac{1}{2}a\,\hat{\mathbf{x}} + \frac{1}{2}b\,\hat{\mathbf{y}}$$
$$\mathbf{a_3} = c\cos\beta\,\hat{\mathbf{x}} + c\sin\beta\,\hat{\mathbf{z}}$$



Basis vectors

		Lattice coordinates		Cartesian coordinates	Wyckoff position	Atom type
$\mathbf{B_1}$	=	0	=	0	(4a)	Ag I
$\mathbf{B_2}$	=	$\frac{1}{2}\,\mathbf{a}_3$	=	$\frac{1}{2}c\cos\beta\,\hat{\mathbf{x}} + \frac{1}{2}c\sin\beta\,\hat{\mathbf{z}}$	(4a)	Ag I
$\mathbf{B_3}$	=	$-y_2\,\mathbf{a}_1 + y_2\,\mathbf{a}_2 + \frac{1}{4}\,\mathbf{a}_3$	=	$\frac{1}{4}c\cos\beta\,\hat{\mathbf{x}} + by_2\,\hat{\mathbf{y}} + \frac{1}{4}c\sin\beta\,\hat{\mathbf{z}}$	(4e)	Ag II
$\mathbf{B_4}$	=	$y_2\,\mathbf{a}_1 - y_2\,\mathbf{a}_2 + \frac{3}{4}\,\mathbf{a}_3$	=	$\frac{3}{4}c\cos\beta\,\hat{\mathbf{x}} - by_2\,\hat{\mathbf{y}} + \frac{3}{4}c\sin\beta\,\hat{\mathbf{z}}$	(4e)	Ag II
$\mathbf{B_5}$	=	$(x_3 - y_3)\,\mathbf{a}_1 + (x_3 + y_3)\,\mathbf{a}_2 + z_3\,\mathbf{a}_3$	=	$(ax_3 + cz_3\cos\beta)\,\hat{\mathbf{x}} + by_3\,\hat{\mathbf{y}} + cz_3\sin\beta\,\hat{\mathbf{z}}$	(8f)	S I
$\mathbf{B_6}$	=	$-(x_3 + y_3)\,\mathbf{a}_1 - (x_3 - y_3)\,\mathbf{a}_2 - \left(z_3 - \frac{1}{2}\right)\,\mathbf{a}_3$	=	$-\left(ax_3 + c\left(z_3 - \frac{1}{2}\right)\cos\beta\right)\,\hat{\mathbf{x}} + by_3\,\hat{\mathbf{y}} - c\left(z_3 - \frac{1}{2}\right)\sin\beta\,\hat{\mathbf{z}}$	(8f)	S I
$\mathbf{B_7}$	=	$-(x_3 - y_3)\,\mathbf{a}_1 - (x_3 + y_3)\,\mathbf{a}_2 - z_3\,\mathbf{a}_3$	=	$-(ax_3 + cz_3\cos\beta)\,\hat{\mathbf{x}} - by_3\,\hat{\mathbf{y}} - cz_3\sin\beta\,\hat{\mathbf{z}}$	(8f)	S I
$\mathbf{B_8}$	=	$(x_3 + y_3)\,\mathbf{a}_1 + (x_3 - y_3)\,\mathbf{a}_2 + \left(z_3 + \frac{1}{2}\right)\,\mathbf{a}_3$	=	$\left(ax_3 + c\left(z_3 + \frac{1}{2}\right)\cos\beta\right)\,\hat{\mathbf{x}} - by_3\,\hat{\mathbf{y}} + c\left(z_3 + \frac{1}{2}\right)\sin\beta\,\hat{\mathbf{z}}$	(8f)	S I
$\mathbf{B_9}$	=	$(x_4 - y_4)\,\mathbf{a}_1 + (x_4 + y_4)\,\mathbf{a}_2 + z_4\,\mathbf{a}_3$	=	$(ax_4 + cz_4\cos\beta)\,\hat{\mathbf{x}} + by_4\,\hat{\mathbf{y}} + cz_4\sin\beta\,\hat{\mathbf{z}}$	(8f)	S II
$\mathbf{B_{10}}$	=	$-(x_4 + y_4)\,\mathbf{a}_1 - (x_4 - y_4)\,\mathbf{a}_2 - \left(z_4 - \frac{1}{2}\right)\,\mathbf{a}_3$	=	$-\left(ax_4 + c\left(z_4 - \frac{1}{2}\right)\cos\beta\right)\,\hat{\mathbf{x}} + by_4\,\hat{\mathbf{y}} - c\left(z_4 - \frac{1}{2}\right)\sin\beta\,\hat{\mathbf{z}}$	(8f)	S II
$\mathbf{B_{11}}$	=	$-(x_4 - y_4)\,\mathbf{a}_1 - (x_4 + y_4)\,\mathbf{a}_2 - z_4\,\mathbf{a}_3$	=	$-(ax_4 + cz_4\cos\beta)\,\hat{\mathbf{x}} - by_4\,\hat{\mathbf{y}} - cz_4\sin\beta\,\hat{\mathbf{z}}$	(8f)	S II
$\mathbf{B_{12}}$	=	$(x_4 + y_4)\,\mathbf{a}_1 + (x_4 - y_4)\,\mathbf{a}_2 + \left(z_4 + \frac{1}{2}\right)\,\mathbf{a}_3$	=	$\left(ax_4 + c\left(z_4 + \frac{1}{2}\right)\cos\beta\right)\,\hat{\mathbf{x}} - by_4\,\hat{\mathbf{y}} + c\left(z_4 + \frac{1}{2}\right)\sin\beta\,\hat{\mathbf{z}}$	(8f)	S II
$\mathbf{B_{13}}$	=	$(x_5 - y_5)\,\mathbf{a}_1 + (x_5 + y_5)\,\mathbf{a}_2 + z_5\,\mathbf{a}_3$	=	$(ax_5 + cz_5\cos\beta)\,\hat{\mathbf{x}} + by_5\,\hat{\mathbf{y}} + cz_5\sin\beta\,\hat{\mathbf{z}}$	(8f)	Sb I
$\mathbf{B_{14}}$	=	$-(x_5 + y_5)\,\mathbf{a}_1 - (x_5 - y_5)\,\mathbf{a}_2 - \left(z_5 - \frac{1}{2}\right)\,\mathbf{a}_3$	=	$-\left(ax_5 + c\left(z_5 - \frac{1}{2}\right)\cos\beta\right)\,\hat{\mathbf{x}} + by_5\,\hat{\mathbf{y}} - c\left(z_5 - \frac{1}{2}\right)\sin\beta\,\hat{\mathbf{z}}$	(8f)	Sb I
$\mathbf{B_{15}}$	=	$-(x_5 - y_5)\,\mathbf{a}_1 - (x_5 + y_5)\,\mathbf{a}_2 - z_5\,\mathbf{a}_3$	=	$-(ax_5 + cz_5\cos\beta)\,\hat{\mathbf{x}} - by_5\,\hat{\mathbf{y}} - cz_5\sin\beta\,\hat{\mathbf{z}}$	(8f)	Sb I
$\mathbf{B_{16}}$	=	$(x_5 + y_5)\,\mathbf{a}_1 + (x_5 - y_5)\,\mathbf{a}_2 + \left(z_5 + \frac{1}{2}\right)\,\mathbf{a}_3$	=	$\left(ax_5 + c\left(z_5 + \frac{1}{2}\right)\cos\beta\right)\,\hat{\mathbf{x}} - by_5\,\hat{\mathbf{y}} + c\left(z_5 + \frac{1}{2}\right)\sin\beta\,\hat{\mathbf{z}}$	(8f)	Sb I

References

[1] H. Effenberger, W. H. Paar, D. Topa, A. J. Criddle, and M. Fleck, *The new mineral baumstarkite and a structural reinvestigation of aramayoite and miargyrite*, Am. Mineral. **87**, 753–764 (2002).

Found in

[1] R. T. Downs and M. Hall-Wallace, *The American Mineralogist Crystal Structure Database*, Am. Mineral. **88**, 247–250 (2003).