


NiO₂ Structure:

AB2_mC6_12_a_i-003

Cite this page as: H. Eckert, S. Divilov, A. Zettel, M. J. Mehl, D. Hicks, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 4*. In preparation.

<https://aflow.org/p/DZY0>

https://aflow.org/p/AB2_mC6_12_a_i-003




Prototype	NiO ₂
AFLOW prototype label	AB2_mC6_12_a_i-003
ICSD	88720
Pearson symbol	mC6
Space group number	12
Space group symbol	<i>C</i> 2/ <i>m</i>
AFLOW prototype command	<code>aflow --proto=AB2_mC6_12_a_i-003 --params=a, b/a, c/a, β, x₂, z₂</code>

- *C*34 Calverite, α-HgO₂, and NiO₂ (this structure) have the same AFLOW prototype label, AB2_mC6_12_a.i. The structures are generated by the same symmetry operations with different sets of parameters (`--params`) specified in their corresponding CIF files.

Base-centered Monoclinic primitive vectors

$$\begin{aligned} \mathbf{a}_1 &= \frac{1}{2}a \hat{\mathbf{x}} - \frac{1}{2}b \hat{\mathbf{y}} \\ \mathbf{a}_2 &= \frac{1}{2}a \hat{\mathbf{x}} + \frac{1}{2}b \hat{\mathbf{y}} \\ \mathbf{a}_3 &= c \cos \beta \hat{\mathbf{x}} + c \sin \beta \hat{\mathbf{z}} \end{aligned}$$



Basis vectors

	Lattice coordinates	=	Cartesian coordinates	Wyckoff position	Atom type
\mathbf{B}_1	=	0	=	0	(2a) Ni I
\mathbf{B}_2	=	$x_2 \mathbf{a}_1 + x_2 \mathbf{a}_2 + z_2 \mathbf{a}_3$	=	$(ax_2 + cz_2 \cos \beta) \hat{\mathbf{x}} + cz_2 \sin \beta \hat{\mathbf{z}}$	(4i) O I
\mathbf{B}_3	=	$-x_2 \mathbf{a}_1 - x_2 \mathbf{a}_2 - z_2 \mathbf{a}_3$	=	$-(ax_2 + cz_2 \cos \beta) \hat{\mathbf{x}} - cz_2 \sin \beta \hat{\mathbf{z}}$	(4i) O I

References

- [1] J. M. Tarascon, G. Vaughan, Y. Chabre, L. Seguin, M. Anne, P. Strobel, and G. Amatucci, In Situ *Structural and Electrochemical Study of $Ni_{1-x}Co_xO_2$ Metastable Oxides Prepared by Soft Chemistry*, J. Solid State Chem. **147**, 410–420 (1999), doi:10.1006/jssc.1999.8465.