$CoGeO_3$ Structure: ABC3_mC40_15_2e_f_3f-001

Cite this page as: H. Eckert, S. Divilov, A. Zettel, M. J. Mehl, D. Hicks, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 4*. In preparation.

https://aflow.org/p/6YLQ

https://aflow.org/p/ABC3_mC40_15_2e_f_3f-001



Prototype	$CoGeO_3$
AFLOW prototype label	ABC3_mC40_15_2e_f_3f-001
ICSD	26814
Pearson symbol	mC40
Space group number	15
Space group symbol	$C2/c$
AFLOW prototype command	`aflow --proto=ABC3_mC40_15_2e_f_3f-001 --params=`$a, b/a, c/a, \beta, y_1, y_2, x_3, y_3, z_3, x_4, y_4, z_4, x_5, y_5, z_5, x_6, y_6, z_6$

Other compounds with this structure
$MgGeO_3$

- This is the ground state structure of $CoGeO_3$. Above 1351°C it transforms to the tetragonal $MgSiO_3$ ($S4_3$) structure, which remains metastable down to 4K (Zhao, 2021).
- The crystal structure data was taken at 4K.
- There is no ICSD or CCDC entry for (Redhammer, 2010). Instead we use the ICSD from the much older work of (Peacor, 1968).

Base-centered Monoclinic primitive vectors

$$
\begin{aligned}
\mathbf{a_1} &= \tfrac{1}{2}a\,\hat{\mathbf{x}} - \tfrac{1}{2}b\,\hat{\mathbf{y}} \\
\mathbf{a_2} &= \tfrac{1}{2}a\,\hat{\mathbf{x}} + \tfrac{1}{2}b\,\hat{\mathbf{y}} \\
\mathbf{a_3} &= c\cos\beta\,\hat{\mathbf{x}} + c\sin\beta\,\hat{\mathbf{z}}
\end{aligned}
$$



Basis vectors

		Lattice coordinates		Cartesian coordinates	Wyckoff position	Atom type
$\mathbf{B_1}$	=	$-y_1\,\mathbf{a}_1 + y_1\,\mathbf{a}_2 + \frac{1}{4}\,\mathbf{a}_3$	=	$\frac{1}{4}c\cos\beta\,\hat{\mathbf{x}} + by_1\,\hat{\mathbf{y}} + \frac{1}{4}c\sin\beta\,\hat{\mathbf{z}}$	(4e)	Co I
$\mathbf{B_2}$	=	$y_1\,\mathbf{a}_1 - y_1\,\mathbf{a}_2 + \frac{3}{4}\,\mathbf{a}_3$	=	$\frac{3}{4}c\cos\beta\,\hat{\mathbf{x}} - by_1\,\hat{\mathbf{y}} + \frac{3}{4}c\sin\beta\,\hat{\mathbf{z}}$	(4e)	Co I
$\mathbf{B_3}$	=	$-y_2\,\mathbf{a}_1 + y_2\,\mathbf{a}_2 + \frac{1}{4}\,\mathbf{a}_3$	=	$\frac{1}{4}c\cos\beta\,\hat{\mathbf{x}} + by_2\,\hat{\mathbf{y}} + \frac{1}{4}c\sin\beta\,\hat{\mathbf{z}}$	(4e)	Co II
$\mathbf{B_4}$	=	$y_2\,\mathbf{a}_1 - y_2\,\mathbf{a}_2 + \frac{3}{4}\,\mathbf{a}_3$	=	$\frac{3}{4}c\cos\beta\,\hat{\mathbf{x}} - by_2\,\hat{\mathbf{y}} + \frac{3}{4}c\sin\beta\,\hat{\mathbf{z}}$	(4e)	Co II
$\mathbf{B_5}$	=	$(x_3 - y_3)\,\mathbf{a}_1 + (x_3 + y_3)\,\mathbf{a}_2 + z_3\,\mathbf{a}_3$	=	$(ax_3 + cz_3\cos\beta)\,\hat{\mathbf{x}} + by_3\,\hat{\mathbf{y}} + cz_3\sin\beta\,\hat{\mathbf{z}}$	(8f)	Ge I
$\mathbf{B_6}$	=	$-(x_3 + y_3)\,\mathbf{a}_1 - (x_3 - y_3)\,\mathbf{a}_2 - \left(z_3 - \frac{1}{2}\right)\,\mathbf{a}_3$	=	$-\left(ax_3 + c\left(z_3 - \frac{1}{2}\right)\cos\beta\right)\,\hat{\mathbf{x}} + by_3\,\hat{\mathbf{y}} - c\left(z_3 - \frac{1}{2}\right)\sin\beta\,\hat{\mathbf{z}}$	(8f)	Ge I
$\mathbf{B_7}$	=	$-(x_3 - y_3)\,\mathbf{a}_1 - (x_3 + y_3)\,\mathbf{a}_2 - z_3\,\mathbf{a}_3$	=	$-(ax_3 + cz_3\cos\beta)\,\hat{\mathbf{x}} - by_3\,\hat{\mathbf{y}} - cz_3\sin\beta\,\hat{\mathbf{z}}$	(8f)	Ge I
$\mathbf{B_8}$	=	$(x_3 + y_3)\,\mathbf{a}_1 + (x_3 - y_3)\,\mathbf{a}_2 + \left(z_3 + \frac{1}{2}\right)\,\mathbf{a}_3$	=	$\left(ax_3 + c\left(z_3 + \frac{1}{2}\right)\cos\beta\right)\,\hat{\mathbf{x}} - by_3\,\hat{\mathbf{y}} + c\left(z_3 + \frac{1}{2}\right)\sin\beta\,\hat{\mathbf{z}}$	(8f)	Ge I
$\mathbf{B_9}$	=	$(x_4 - y_4)\,\mathbf{a}_1 + (x_4 + y_4)\,\mathbf{a}_2 + z_4\,\mathbf{a}_3$	=	$(ax_4 + cz_4\cos\beta)\,\hat{\mathbf{x}} + by_4\,\hat{\mathbf{y}} + cz_4\sin\beta\,\hat{\mathbf{z}}$	(8f)	O I
$\mathbf{B_{10}}$	=	$-(x_4 + y_4)\,\mathbf{a}_1 - (x_4 - y_4)\,\mathbf{a}_2 - \left(z_4 - \frac{1}{2}\right)\,\mathbf{a}_3$	=	$-\left(ax_4 + c\left(z_4 - \frac{1}{2}\right)\cos\beta\right)\,\hat{\mathbf{x}} + by_4\,\hat{\mathbf{y}} - c\left(z_4 - \frac{1}{2}\right)\sin\beta\,\hat{\mathbf{z}}$	(8f)	O I
$\mathbf{B_{11}}$	=	$-(x_4 - y_4)\,\mathbf{a}_1 - (x_4 + y_4)\,\mathbf{a}_2 - z_4\,\mathbf{a}_3$	=	$-(ax_4 + cz_4\cos\beta)\,\hat{\mathbf{x}} - by_4\,\hat{\mathbf{y}} - cz_4\sin\beta\,\hat{\mathbf{z}}$	(8f)	O I
$\mathbf{B_{12}}$	=	$(x_4 + y_4)\,\mathbf{a}_1 + (x_4 - y_4)\,\mathbf{a}_2 + \left(z_4 + \frac{1}{2}\right)\,\mathbf{a}_3$	=	$\left(ax_4 + c\left(z_4 + \frac{1}{2}\right)\cos\beta\right)\,\hat{\mathbf{x}} - by_4\,\hat{\mathbf{y}} + c\left(z_4 + \frac{1}{2}\right)\sin\beta\,\hat{\mathbf{z}}$	(8f)	O I
$\mathbf{B_{13}}$	=	$(x_5 - y_5)\,\mathbf{a}_1 + (x_5 + y_5)\,\mathbf{a}_2 + z_5\,\mathbf{a}_3$	=	$(ax_5 + cz_5\cos\beta)\,\hat{\mathbf{x}} + by_5\,\hat{\mathbf{y}} + cz_5\sin\beta\,\hat{\mathbf{z}}$	(8f)	O II
$\mathbf{B_{14}}$	=	$-(x_5 + y_5)\,\mathbf{a}_1 - (x_5 - y_5)\,\mathbf{a}_2 - \left(z_5 - \frac{1}{2}\right)\,\mathbf{a}_3$	=	$-\left(ax_5 + c\left(z_5 - \frac{1}{2}\right)\cos\beta\right)\,\hat{\mathbf{x}} + by_5\,\hat{\mathbf{y}} - c\left(z_5 - \frac{1}{2}\right)\sin\beta\,\hat{\mathbf{z}}$	(8f)	O II
$\mathbf{B_{15}}$	=	$-(x_5 - y_5)\,\mathbf{a}_1 - (x_5 + y_5)\,\mathbf{a}_2 - z_5\,\mathbf{a}_3$	=	$-(ax_5 + cz_5\cos\beta)\,\hat{\mathbf{x}} - by_5\,\hat{\mathbf{y}} - cz_5\sin\beta\,\hat{\mathbf{z}}$	(8f)	O II
$\mathbf{B_{16}}$	=	$(x_5 + y_5)\,\mathbf{a}_1 + (x_5 - y_5)\,\mathbf{a}_2 + \left(z_5 + \frac{1}{2}\right)\,\mathbf{a}_3$	=	$\left(ax_5 + c\left(z_5 + \frac{1}{2}\right)\cos\beta\right)\,\hat{\mathbf{x}} - by_5\,\hat{\mathbf{y}} + c\left(z_5 + \frac{1}{2}\right)\sin\beta\,\hat{\mathbf{z}}$	(8f)	O II
$\mathbf{B_{17}}$	=	$(x_6 - y_6)\,\mathbf{a}_1 + (x_6 + y_6)\,\mathbf{a}_2 + z_6\,\mathbf{a}_3$	=	$(ax_6 + cz_6\cos\beta)\,\hat{\mathbf{x}} + by_6\,\hat{\mathbf{y}} + cz_6\sin\beta\,\hat{\mathbf{z}}$	(8f)	O III
$\mathbf{B_{18}}$	=	$-(x_6 + y_6)\,\mathbf{a}_1 - (x_6 - y_6)\,\mathbf{a}_2 - \left(z_6 - \frac{1}{2}\right)\,\mathbf{a}_3$	=	$-\left(ax_6 + c\left(z_6 - \frac{1}{2}\right)\cos\beta\right)\,\hat{\mathbf{x}} + by_6\,\hat{\mathbf{y}} - c\left(z_6 - \frac{1}{2}\right)\sin\beta\,\hat{\mathbf{z}}$	(8f)	O III

$\mathbf{B_{19}}$	=	$-(x_6 - y_6)\ \mathbf{a}_1 - (x_6 + y_6)\ \mathbf{a}_2 - z_6\ \mathbf{a}_3$	=	$-(ax_6 + cz_6 \cos\beta)\ \hat{\mathbf{x}} - by_6\ \hat{\mathbf{y}} - cz_6 \sin\beta\ \hat{\mathbf{z}}$	(8f)	O III
$\mathbf{B_{20}}$	=	$(x_6 + y_6)\ \mathbf{a}_1 + (x_6 - y_6)\ \mathbf{a}_2 + \left(z_6 + \frac{1}{2}\right)\ \mathbf{a}_3$	=	$\left(ax_6 + c\left(z_6 + \frac{1}{2}\right)\cos\beta\right)\ \hat{\mathbf{x}} - by_6\ \hat{\mathbf{y}} + c\left(z_6 + \frac{1}{2}\right)\sin\beta\ \hat{\mathbf{z}}$	(8f)	O III

References

[1] G. J. Redhammer, A. Senyshyn, G. Tippelt, C. Pietzonka, G. Roth, and G. Amthauer, *Magnetic and nuclear structure and thermal expansion of orthorhombic and monoclinic polymorphs of $CoGeO_3$ pyroxene*, Phys. Chem. Minerals **37**, 311–332 (2010), doi:10.1007/s00269-009-0335-x.

[2] D. R. Peacor, *The crystal structure of $CoGeO_3$*, Z. Krystallogr. **126**, 299–306 (1968), doi:10.1524/zkri.1968.126.16.299.