

InS Structure:

AB_oP8_58_g-g-001





This structure originally had the label AB_oP8_58_g-g. Calls to that address will be redirected here.

Cite this page as: D. Hicks, M. J. Mehl, M. Esters, C. Osse, O. Levy, G. L. W. Hart, C. Toher, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 3*, Comput. Mater. Sci. **199**, 110450 (2021), doi: 10.1016/j.commatsci.2021.110450.

<https://aflow.org/p/A7M6>

https://aflow.org/p/AB_oP8_58_g-g-001

■ In
■ S



Prototype

InS

AFLOW prototype label

AB_oP8_58_g-g-001

ICSD

15931

Pearson symbol

oP8

Space group number

58

Space group symbol

$Pnnm$

AFLOW prototype command

aflow --proto=AB_oP8_58_g-g-001
--params=a, b/a, c/a, x1, y1, x2, y2


Simple Orthorhombic primitive vectors

$$\mathbf{a}_1 = a \hat{x}$$

$$\mathbf{a}_2 = b \hat{y}$$

$$\mathbf{a}_3 = c \hat{z}$$

$$\begin{matrix} \mathbf{a}_3 \\ \mathbf{a}_1 \\ \mathbf{a}_2 \end{matrix}$$



Basis vectors

	Lattice coordinates		Cartesian coordinates	Wyckoff position	Atom type
B₁	= $x_1 \mathbf{a}_1 + y_1 \mathbf{a}_2$	=	$ax_1 \hat{\mathbf{x}} + by_1 \hat{\mathbf{y}}$	(4g)	In I
B₂	= $-x_1 \mathbf{a}_1 - y_1 \mathbf{a}_2$	=	$-ax_1 \hat{\mathbf{x}} - by_1 \hat{\mathbf{y}}$	(4g)	In I
B₃	= $-\left(x_1 - \frac{1}{2}\right) \mathbf{a}_1 + \left(y_1 + \frac{1}{2}\right) \mathbf{a}_2 + \frac{1}{2} \mathbf{a}_3$	=	$-a\left(x_1 - \frac{1}{2}\right) \hat{\mathbf{x}} + b\left(y_1 + \frac{1}{2}\right) \hat{\mathbf{y}} + \frac{1}{2}c \hat{\mathbf{z}}$	(4g)	In I
B₄	= $\left(x_1 + \frac{1}{2}\right) \mathbf{a}_1 - \left(y_1 - \frac{1}{2}\right) \mathbf{a}_2 + \frac{1}{2} \mathbf{a}_3$	=	$a\left(x_1 + \frac{1}{2}\right) \hat{\mathbf{x}} - b\left(y_1 - \frac{1}{2}\right) \hat{\mathbf{y}} + \frac{1}{2}c \hat{\mathbf{z}}$	(4g)	In I
B₅	= $x_2 \mathbf{a}_1 + y_2 \mathbf{a}_2$	=	$ax_2 \hat{\mathbf{x}} + by_2 \hat{\mathbf{y}}$	(4g)	S I
B₆	= $-x_2 \mathbf{a}_1 - y_2 \mathbf{a}_2$	=	$-ax_2 \hat{\mathbf{x}} - by_2 \hat{\mathbf{y}}$	(4g)	S I
B₇	= $-\left(x_2 - \frac{1}{2}\right) \mathbf{a}_1 + \left(y_2 + \frac{1}{2}\right) \mathbf{a}_2 + \frac{1}{2} \mathbf{a}_3$	=	$-a\left(x_2 - \frac{1}{2}\right) \hat{\mathbf{x}} + b\left(y_2 + \frac{1}{2}\right) \hat{\mathbf{y}} + \frac{1}{2}c \hat{\mathbf{z}}$	(4g)	S I
B₈	= $\left(x_2 + \frac{1}{2}\right) \mathbf{a}_1 - \left(y_2 - \frac{1}{2}\right) \mathbf{a}_2 + \frac{1}{2} \mathbf{a}_3$	=	$a\left(x_2 + \frac{1}{2}\right) \hat{\mathbf{x}} - b\left(y_2 - \frac{1}{2}\right) \hat{\mathbf{y}} + \frac{1}{2}c \hat{\mathbf{z}}$	(4g)	S I

References

- [1] K. Schubert, E. Dörre, and E. Günzel, *Kristallchemische Ergebnisse an Phasen aus B-Elementen*, Naturwissenschaften **41**, 448 (1954), doi:10.1007/BF00628872.

Found in

- [1] P. Villars and L. Calvert, *Pearson's Handbook of Crystallographic Data for Intermetallic Phases* (ASM International, Materials Park, OH, 1991), 2nd edn.